

Opportunities and Challenges for Dietary Arsenic Intervention

Keeve E. Nachman,^{1,2,3,4} Tracy Punshon,^{5,6,7} Laurie Rardin,⁶ Antonio J. Signes-Pastor,^{6,7,9} Carolyn J. Murray,⁷ Brian P. Jackson,^{6,8} Mary Lou Guerinot,⁵ Thomas A. Burke,^{1,3} Celia Y. Chen,^{5,6} Habibul Ahsan,¹⁰ Maria Argos,¹¹ Kathryn L. Cottingham,^{5,7} Francesco Cubadda,¹² Gary L. Ginsberg,¹³ Britton C. Goodale,^{6,14} Margaret Kurzius-Spencer,^{15,16} Andrew A. Meharg,¹⁷ Mark D. Miller,¹⁸ Anne E. Nigra,¹⁹ Claire B. Pendergrast,²⁰ Andrea Raab,²¹ Ken Reimer,²² Kirk G. Scheckel,²³ Tanja Schwerdtle,²⁴ Vivien F. Taylor,^{6,8} Erik J. Tokar,²⁵ Todd M. Warczak,⁵ and Margaret R. Karagas^{6,7,9}

¹Risk Sciences and Public Policy Institute, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland, USA

²Department of Environmental Health and Engineering, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland, USA

³Department of Health Policy and Management, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland, USA

⁴Johns Hopkins Center for a Livable Future, Johns Hopkins University, Baltimore, Maryland, USA

⁵Department of Biological Sciences, Dartmouth College, Hanover, New Hampshire, USA

⁶Dartmouth Superfund Research Program, Hanover, New Hampshire, USA

⁷Dartmouth Children's Environmental Health and Disease Prevention Research Center, Geisel School of Medicine at Dartmouth, Lebanon, New Hampshire, USA

⁸Department of Earth Sciences, Dartmouth College, Hanover, New Hampshire, USA

⁹Department of Epidemiology, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA

¹⁰Department of Public Health Sciences, University of Chicago, Chicago, Illinois, USA

¹¹Division of Epidemiology and Biostatistics, School of Public Health, University of Illinois at Chicago, Chicago, Illinois, USA

¹²Department of Food Safety, Nutrition and Veterinary Public Health, Istituto Superiore di Sanità – Italian National Institute of Health, Rome, Italy

¹³Yale School of Public Health, 60 College St, New Haven, Connecticut, USA

¹⁴Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA

¹⁵Department of Pediatrics, College of Medicine, University of Arizona, Tucson, Arizona, USA

¹⁶Department of Community, Environment and Policy, Mel & Enid College of Public Health, University of Arizona, Tucson, Arizona, USA

¹⁷Institute for Global Food Security, Queen's University Belfast, David Keir Building, Malone Road, Belfast, BT9 5BN, Northern Ireland, UK

¹⁸Western States Pediatric Environmental Health Specialty Unit, University of California, San Francisco, San Francisco, California, USA

¹⁹Department of Environmental Health Sciences, Columbia University Mailman School of Public Health, New York, New York, USA

²⁰University of Washington School of Public Health, Seattle, Washington, USA

²¹Department of Chemistry, University of Aberdeen, Aberdeen, UK

²²Royal Military College, Kingston, Ontario, Canada

²³Land and Materials Management Division, National Risk Management Research Laboratory, United States Environmental Protection Agency, Cincinnati, Ohio, USA

²⁴Institute of Nutritional Sciences, University of Potsdam, Germany

²⁵National Toxicology Program Laboratory, National Toxicology Program, National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina, USA

SUMMARY: The diet is emerging as the dominant source of arsenic exposure for most of the U.S. population. Despite this, limited regulatory efforts have been aimed at mitigating exposure, and the role of diet in arsenic exposure and disease processes remains understudied. In this brief, we discuss the evidence linking dietary arsenic intake to human disease and discuss challenges associated with exposure characterization and efforts to quantify risks. In light of these challenges, and in recognition of the potential longer-term process of establishing regulation, we introduce a framework for shorter-term interventions that employs a field-to-plate food supply chain model to identify monitoring, intervention, and communication opportunities as part of a multisector, multiagency, science-informed, public health systems approach to mitigation of dietary arsenic exposure. Such an approach is dependent on coordination across commodity producers, the food industry, nongovernmental organizations, health professionals, researchers, and the regulatory community. <https://doi.org/10.1289/EHP3997>

Background

Emerging data suggest that arsenic, even at relatively low levels of exposure, may affect human health, particularly during early life (NRC 2014). In consideration of this issue, regulatory efforts have been mobilized to address exposures, primarily from drinking water (Nachman et al. 2017). As successes are increasingly documented in these exposure-reduction efforts (Nigra et al. 2017; Welch et al. 2018), we view the diet as a driving source of exposure in populations with low drinking water arsenic. Despite

its contributions to aggregate arsenic exposures, efforts to tackle arsenic in food have been relatively sparse. The Collaborative on Food with Arsenic and associated Risk and Regulation (C-FARR) brought arsenic and food scientists together with policy stakeholders for a workshop focusing on knowledge gaps and policy questions in recognition of this lagging focus. The resulting five papers address this issue from soil to plate to policy (Cubadda et al. 2017; Davis et al. 2017; Nachman et al. 2017; Punshon et al. 2017; Taylor et al. 2017). Moving beyond the C-FARR workshop, and considering scientific and policy hurdles, we discuss here an array of immediate-term opportunities that exist among multiple stakeholder groups to intervene on dietary arsenic exposures.

Arsenic exposure through drinking water has established health impacts (NRC 2014), and although existing science supports the assumption that effects will be similar from food, the role of diet in population arsenic exposures has been less studied. Most epidemiologic studies describing arsenic's effects have relied on drinking water concentrations or urine measurements as exposure measures, which are not designed to disentangle the role of dietary arsenic in the occurrence of disease (Nachman et al. 2017). Further, in comparison with drinking water, where arsenic occurs solely in inorganic forms, the species profile for

Address correspondence to K.E. Nachman, 615 N. Wolfe St., W7010-E, Baltimore, MD 21205, USA. Telephone: (410) 223-1811; Fax: (410) 502-7579. Email: knachman@jhu.edu

The authors declare they have no actual or potential competing financial interests.

Received 3 June 2018; Revised 16 July 2018; Accepted 20 July 2018; Published 31 August 2018.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

food is complex, comprising inorganic arsenic (arsenite and arsenate) and more than 100 organic arsenic species that vary within and across foods (Cubadda et al. 2017). Although epidemiologic studies have provided clear evidence for the carcinogenicity of inorganic arsenic (iAs) and link it to myriad noncancer outcomes, our understanding of the chronic toxicity of organic species is emerging, further complicating characterization of dietary arsenic exposures (Taylor et al. 2017).

Although drinking water may be an easier medium for epidemiologists to use to examine arsenic exposure, it is not the dominant pathway for most of the U.S. population. An earlier assessment of public water in 25 states estimated that <4% of people were served by systems with arsenic concentrations >1 µg/L, and just over 1% were above the 10 µg/L U.S. EPA Maximum Contaminant Level (MCL) (Mushak et al. 2000). Since then, public water exposures have declined, following the implementation of the MCL (Nigra et al. 2017). Evaluations of modeled exposure across dietary and drinking water pathways suggest that in those with drinking water arsenic concentrations below the MCL, diet is responsible for the majority of both total and iAs intake (Kurzius-Spencer et al. 2014; Xue et al. 2010).

Despite the importance of the dietary pathway, sparse data exist about the role of dietary arsenic exposure in disease processes. Among these investigations, most have focused on rice, and very few have jointly considered drinking water exposures along with other potential dietary sources. Additional challenges in characterizing the role of diet include difficulty in separating out the effects of arsenic in the water used to cook rice in high water arsenic regions, estimating arsenic levels in rice and rice products and adjusting for the effects of dietary and other confounders (e.g., carbohydrate content of rice). To date, few studies raise the possibility of adverse health impacts of rice consumption among those with relatively low drinking water arsenic concentrations. These studies found a trend in cardiovascular disease risk with white (but not brown or overall) rice consumption in the prospective Nurses' Health Study among participants living in low arsenic regions (<3 µg/L), but not among those living in higher arsenic regions (Muraki et al. 2015). In Bangladesh, Melkonian et al. (2013) found a relationship between steamed rice consumption and premalignant and malignant skin lesions in both cross-sectional and prospective analyses of the Health Effects of Arsenic Longitudinal Study only among those with drinking water arsenic concentrations <100 µg/L. In the United States, an association between rice consumption and squamous cell carcinoma of the skin was observed in a population-based study, largely among those with tap water concentrations of arsenic <1 µg/L (Gossai et al. 2017). Using data from the Nurses' Health Study and the Health Professionals Follow-up Study, Zhang et al. (2016) found no significant increases in cancer risk from rice consumption but identified a borderline significant increased bladder cancer risk when comparing the highest rice-consumption group to the lowest; the study did not involve consideration of drinking water arsenic. Thus, although limited, epidemiologic evidence currently suggests that the effects of rice consumption may mirror those of arsenic in drinking water.

Dietary exposure characterization poses unique methodological challenges, as few studies describe distributions of arsenic across foods, and far fewer involve speciation (Cubadda et al. 2017). These data gaps complicate efforts to quantify dietary exposures and limit the ability to target foods for intervention based on their relative contributions to aggregate intake.

Though risks from food cannot be precisely quantified, existing evidence supports efforts to limit exposure wherever possible, especially for vulnerable populations (Naujokas et al. 2013). Despite the dominance of diet in population arsenic

exposures, arsenic in food remains unregulated in the United States (Nachman et al. 2017). Under normal circumstances, regulatory agencies would be expected to play a key role in drafting enforceable interventions aimed at minimizing population exposures. In the case of arsenic, no enforceable regulatory standards exist for any foods, though the U.S. Food and Drug Administration (U.S. FDA) has proposed draft action levels and accompanying industry guidance for apple juice (U.S. FDA 2013) and infant rice cereal (U.S. FDA 2016). Legislative attempts to force stringent regulatory action (U.S. Congress 2012, 2015, 2017) have thus far been unsuccessful, though in its 2018 evaluation of U.S. FDA and U.S. Department of Agriculture (USDA) action on arsenic in rice, the U.S. Government Accountability Office (U.S. GAO) recommended FDA finalize its guidance on arsenic in infant rice cereal and called for improvements in interagency coordination on analytical methods and risk-assessment approaches for arsenic and other food contaminants (U.S. GAO 2018).

We previously described an approach that could be used to reduce dietary iAs exposure in the U.S. population. It employs existing monitoring data (and proposes newer data streams) to facilitate an iterative reevaluation of population dietary iAs exposures, and subsequently prioritizes specific foods for intervention, both in the form of measures to reduce arsenic concentrations in key foods (via action levels and regulatory standards to compel producer efforts to reduce arsenic content) and dietary advice when those reductions are more difficult (Nachman et al. 2017). If adequately protective standards are implemented and properly enforced (Signes-Pastor et al. 2017b), this type of holistic approach, which recognizes the pervasiveness of iAs in the food supply (Schoof et al. 1999), may hold promise for reducing population exposures. To complement this approach, especially in circumstances where regulatory action may occur more slowly (Krisberg 2017; Samet et al. 2017), other opportunities exist that can be implemented in the short term and may yield public health benefits stemming from arsenic exposure reductions.

Discussion

In the absence of regulations specific to arsenic in food, we argue that there are many opportunities for nonregulatory stakeholders to mitigate dietary exposures based on our current understanding of the food supply chain. Furthermore, we believe it is possible to reduce exposure to arsenic through control of source releases, and demonstration of ways to reduce the arsenic content of retail foods may pave the way for future regulatory changes at the federal level.

Figure 1 depicts intervention and monitoring points along the food supply that may be meaningfully employed by various stakeholders to reduce population dietary exposures. Given that rice has been the subject of considerable intervention research, we use it in demonstrating opportunities. We recognize, however, as we have previously specified (Nachman et al. 2017), that a coordinated, iterative monitoring intervention strategy that considers the contribution of all foods and beverages would begin to address the current needs of reducing exposure.

Rice cultivar selection and agronomic practices dominate as driving factors of grain arsenic concentrations (Norton et al. 2012; Yang et al. 2017). The influence of genetic variation on grain arsenic indicates that breeding low-arsenic rice cultivars is a viable approach and is currently underway (Norton et al. 2009). Manipulating flooding cycles in rice production, tied to sustainable agricultural practices for reducing water use, can also reduce grain arsenic (Li et al. 2009; Yang et al. 2017). Fertilization strategies, particularly using silicon, have potential as an effective strategy to reduce arsenic, and prevent certain plant diseases (e.g., straighthead disease in rice) (Limmer et al. 2018). Equally




Figure 1. The food supply chain and opportunities for dietary arsenic exposure monitoring and intervention. A field-to-plate food supply chain model can aid in identifying monitoring, intervention and communication opportunities as part of a multisector, multiagency, science-informed, public health systems approach to mitigation of dietary arsenic exposure.

important to demonstrating efficacy of these approaches, is making them accessible and attractive to small scale rural farming operations; for example, re-using rice husk as a silicon fertilizer for rice paddy soils [shown to lower total and iAs (Penido et al. 2016; Seyfferth et al. 2016; Teasley et al. 2017)] has great potential because silicon fertilizers are too expensive to be implemented at many smallholding farms, and, if used, would potentially drive up the price of rice.

Recent Codex Alimentarius international food standards codes of practice (Codex 2017) lay out logical steps for the prevention and reduction of arsenic contamination in rice, grouped into: source-directed measures (determining that the arsenic concentrations of soils and irrigation water are not elevated before use; cognizance of proximity to point sources of arsenic contamination; avoiding use of arsenical pesticides and veterinary drugs, feed, soil amendments, and fertilizer), agricultural measures (use of alternative intermittent flooding strategies and low-arsenic cultivars), and monitoring and risk communication programs. Adherence to this code has the potential to dramatically reduce arsenic concentrations, particularly if food-product labeling were used to raise consumer awareness.

Recognizing that the arsenic content of many finished foods is dependent on the raw commodities that serve as ingredients, opportunities exist regarding both the selection of ingredients and quality assurance measures related to the arsenic concentrations of those constituents. For example, the replacement of high-fructose corn syrup with organic brown-rice syrup in products like toddler formula and energy bars (Jackson et al. 2012) and the use of the algae-derived gelatin substitute carrageenan in a variety of foods (Díaz et al. 2012; U.S. FDA 2017b) may contribute to dietary arsenic intake, although the health impacts of constituents require further testing. Currently, no programs exist to monitor the arsenic content of additives or ingredients, and recommended international standards for some of these inputs may not be based upon the most recent evidence (Food and Agriculture Organization 2001). Processor-level preventive actions, such as ingredient monitoring or ingredient substitution with others less likely to accumulate arsenic, may be beneficial.

Additional reductions may be achievable during food and ingredient processing by taking advantage of our understanding of

the nature of iAs accumulation within commodities. For example, rice-processing procedures designed to remove the husk (which has been shown to accumulate iAs at levels as high as 1 ppm) and further polish the grain can achieve substantial iAs reductions (Carey et al. 2015; Meharg et al. 2008; Signes-Pastor et al. 2017a; Signes et al. 2008). Further, lower iAs content in rice-based products can be achieved by dilution of rice with other gluten-free grain (Carey et al. 2018).

In some instances when adequate exposure reduction cannot be accomplished by lowering the arsenic content of foods, alternative approaches to reduce dietary exposures may be warranted, such as modifying consumer preferences to influence purchasing behaviors (which would exert pressure on producers to address the arsenic content of their foods) (Nachman et al. 2017).

Interventions to communicate information on arsenic in food are somewhat limited and often rice-centric, though some helpful sources exist, including a comprehensive website focused on arsenic in rice and rice products by Consumer Reports (2015). The U.S. FDA (2017a) also maintains a communications and data-sharing website for arsenic in food. Research-based consumer education resources highlighting arsenic health risks and dietary (and other) exposure-reduction options are available online in a comprehensive website (Dartmouth College Superfund Research Program 2017) and an interactive graphic (Children's Environmental Health and Disease Prevention Research Center at Dartmouth 2016). Beyond these vetted resources, social media sites and blogs from gluten-free dietary information resources and Celiac disease organizations (e. g., Celiac Disease Foundation 2017) also provide exposure-reduction resources. Research is needed to determine the effectiveness of these tools on individual behavior change and consumer decision-making. Additional communication efforts need to be tailored and targeted to populations at higher risk of adverse health effects from arsenic exposure, such as infants, children, and pregnant women, as well as to populations more highly exposed due to cultural preference and dietary restrictions (Lai et al. 2015).

Communication efforts successful in changing consumer behaviors may have collateral benefits along the food supply chain. Research published by Jackson et al. (2012) identifying elevated iAs in organic brown-rice syrup used in toddler formula resulted in a media cycle and public outcry that prompted some

food manufacturers to take action to address arsenic in their products (Lundberg Family Farms 2016; Nature's One 2018), illustrating how media attention and consumer opinion can drive industry action to reduce arsenic content in food products even without regulation.

Recent research has highlighted post-retail opportunities for consumers to reduce the arsenic content of finished foods. Several studies have shown that some of the iAs content is readily leached from rice by washing and soaking, and even more can be removed by increasing the volume of cooking water that rice encounters, followed by draining off the excess iAs-containing water generated during cooking (Raab et al. 2009; Sengupta et al. 2006; Signes et al. 2008). To this end, a new rice cooking approach has been proposed based on a continual stream of percolating near-boiling water; when low-arsenic water has been used, this method has been demonstrated to remove ~57% of iAs from cooked rice at water-to-rice cooking ratio of 12:1, and a removal of ~70% of iAs with a ratio of 150:1 or higher in cooked rice bran (Signes-Pastor et al. 2017a). Proponents of this approach have suggested its feasibility both in industrial and domestic settings, and possibly even when low arsenic cooking water is not available by percolating freshly recycled distilled water (Carey et al. 2015). Such an approach may provide a near-term, suitable solution to mitigate high iAs levels in rice and rice bran, though further research is needed to assess potential loss of key vitamins and other water-soluble compounds that could be refortified if necessary.

Existing population-level health surveys, especially those that describe dietary patterns and collect biomarker data, can help target dietary interventions, and to a lesser extent, assist in evaluating existing exposure-reduction efforts. The What We Eat in America food survey, conducted as part of the National Health and Nutrition Examination Survey (USDA 2018), enables the characterization of dietary patterns across key population subgroups, including both vulnerable life stages regarding arsenic exposure and population subgroups with higher consumption of arsenic-accumulating foods. Urinary arsenic concentrations have been consistently related to rice consumption among populations throughout the world, and in both interventional and observational studies (Davis et al. 2017). These studies encompass populations of various ages, ranging from infancy and early childhood to pregnancy and later adulthood. Arsenic is known to pass through the placenta (e.g., Hall et al. 2007); in a prospective cohort study, intake of rice during pregnancy was related to arsenic concentration in infant toenails collected within a few months of birth (Davis et al. 2014), and a longitudinal study found an increase of infants' urinary arsenic during transition to solid food, including rice cereals, fruits, and vegetables (Signes-Pastor et al. 2018). Evaluations of arsenic biomarker trends among these subpopulations may allow for identification of interventions aimed at specific subpopulations or foods. Further, recent efforts to disentangle dietary from drinking water exposures within biomarker measurements may enhance sensitivity of evaluation of dietary interventions (Jones et al. 2016).

Conclusions

Dietary arsenic exposures are an increasingly recognized public health concern, and this concern calls for interventions to reduce exposure, especially to vulnerable populations, such as pregnant women, infants, and children. Regulatory advances will be essential for effective intervention to lower dietary arsenic exposure, but opportunities also exist for various stakeholders, including commodity producers, the food industry, physicians, and the public, to take steps along the supply chain. For instance, we have proposed a proactive scheme designed to compel the production

of low-arsenic foods (Nachman et al. 2017). Such approaches require better monitoring efforts, such as those implemented in the Italian Total Diet Study (Cubadda et al. 2016), that employ the latest measurement techniques and push the field of practice forward. Moving further upstream, public health benefits may be derived from policies aimed at creating producer recommendations or setting production standards for soils and irrigation water.

Building upon GAO's (2018) recommendations of inter-agency coordination on arsenic in food, future regulatory policy interventions must consider the relative contributions of different dietary and other pathways to aggregate arsenic exposures. Further, enhanced efforts to monetize the avoidance of non-cancer health outcomes (e.g., cardiovascular, developmental, and many others) associated with arsenic exposure would be of great value. The effect of reducing the occurrence of these outcomes was not quantitatively considered in the U.S. EPA's economic justification of the arsenic in drinking water rule (U.S. EPA 2000). An economic understanding of the public health benefits of preventing these outcomes would likely justify more stringent measures to reduce exposure.

An urgent component of moving regulatory policy forward is updating the iAs toxicological assessment. The EPA IRIS's endeavor to update its iAs risk assessment is currently underway but has yet to be finalized. GAO called for U.S. FDA to update its rice risk assessment (U.S. GAO 2018), however, such efforts may be redundant with, and less comprehensive than that of the EPA IRIS program. In its 2018 review of updates to the IRIS program, the National Research Council (NRC), expressed satisfaction with the program's adoption of systematic review methods in pursuit of hazard characterization and dose-response assessment (NRC 2018). With much effort already invested in the development of the iAs assessment, and ongoing coordination with the NRC, it is anticipated that the toxicity values produced by this effort will be key in informing future FDA rice (and other food) risk assessment endeavors. With these toxicity values, interventions will be able to set sights on risk-based exposure limits, rather than limiting efforts to general exposure reduction in the absence of any target based on actual dose-response relationships.

Certain gaps in understanding of the science exist that, if filled, would bolster policy efforts aimed at exposure reduction. Specifically, primary research, as well as systematic reviews and meta-analyses of "other priority outcomes" (e.g., diabetes, neurodevelopmental toxicity and immune effects) and "other endpoints to consider" (e.g., renal disease, liver and pancreatic cancer, and hypertension), as specified by the NRC (2014), are needed. In addition, research to better characterize life stage-specific windows of vulnerability to arsenic exposure, as well as their interactions with genetic factors, epigenetic alterations, and other contaminant exposures (Cardenas et al. 2015), as well as the timing of gene-expression changes (Wright and Christiani 2010) is warranted. Further, broadening consideration of arsenic to include oral exposure to organic arsenic species commonly found in foods (including methylated forms, arsenosugars, and arsenolipids) would be of value, especially for understanding dietary exposures. Focusing interventions narrowly on inorganic species may not account for the true health burdens resulting from exposure. Although uncertainty remains over population-wide health effects of dietary arsenic, focus on susceptible populations is critical. The mitigating role of nutrients involved in arsenic metabolism, such as folate and other B vitamins (Kurzius-Spencer et al. 2017; Spratlen et al. 2017) strongly suggests that susceptible populations will also include those with gastrointestinal disorders that inhibit the intestinal absorption of these vitamins, such as individuals with Celiac

disease, a population with increased dietary arsenic exposure from the prevalence of a rice-based, gluten-free diet (Bulka et al. 2017; Punshon and Jackson 2018).

Dietary arsenic exposure is an important public health challenge that contributes to population risk of a broad number of adverse health impacts. Current regulatory approaches for arsenic are limited, and the statutory basis for potential controls is fragmented across multiple agencies. As a result, the patchwork of state and federal efforts has had limited success in reducing dietary exposures and often results in mixed messages to consumers about risks. It is time to rethink our current approaches and develop a systems approach to dietary arsenic exposures — from source to ingestion. A renewed, inclusive approach to problem formulation, including the social and behavioral sciences (The National Academies 2018), can provide a renewed perspective on both efforts to characterize the scope of the problem and the most productive opportunities to intervene. Our greatest success in reducing dietary exposures can be achieved only by a multisector, multiagency, science-informed, public health systems approach coordinated across regulators and the industry (Burke et al. 2017). When those are not enough to ensure minimal arsenic contributions through diet, a well-informed public empowered to make the right food choices is essential.

Acknowledgments

This paper, a product of the Collaborative on Food with Arsenic and associated Risk and Regulation (C-FARR), is supported by the Dartmouth College Toxic Metals Superfund Research Program through funds from the National Institute of Environmental Health Sciences (NIEHS) of the National Institutes of Health (NIH) under Award Number 1R13ES026493-01 to C. C. and Award Number P42ES007373 to Bruce Stanton, and the Children's Environmental Health and Disease Prevention Research Center at Dartmouth through funds from the NIEHS of the NIH under Award Number P01ES022832 and from the U.S. Environmental Protection Agency RD-83544201 to M. K. The views expressed in this paper are the those of the authors and do not necessarily reflect the official views of any agency of the United States or other government. We are grateful to A. Seyffert for helpful discussions regarding silica and rice production.

References

Bulka CM, Davis MA, Karagas MR, Ahsan H, Argos M. 2017. The unintended consequences of a gluten-free diet. *Epidemiology* 28(3):e24–e25, PMID: 28166100, <https://doi.org/10.1097/EDE.0000000000000640>.

Burke TA, Cascio WE, Costa DL, Deener K, Fontaine TD, Fulk FA, et al. 2017. Rethinking environmental protection: meeting the challenges of a changing world. *Environ Health Perspect* 125(3):A43–A49, PMID: 28248180, <https://doi.org/10.1289/EHP1465>.

Cardenas A, Koestler DC, Houseman EA, Jackson BP, Kile ML, Karagas MR, et al. 2015. Differential DNA methylation in umbilical cord blood of infants exposed to mercury and arsenic in utero. *Epigenetics* 10(6):508–515, PMID: 25923418, <https://doi.org/10.1080/15592294.2015.1046026>.

Carey M, Jiujiun X, Farias JG, Meharg AA. 2015. Rethinking rice preparation for highly efficient removal of inorganic arsenic using percolating cooking water. *PLoS One* 10(7):e0131608, PMID: 26200355, <https://doi.org/10.1371/journal.pone.0131608>.

Carey M, Donaldson E, Signes-Pastor AJ, Meharg AA. 2018. Dilution of rice with other gluten free grains to lower inorganic arsenic in foods for young children in response to European Union regulations provides impetus to setting stricter standards. *PLoS One* 13(3):e0194700, PMID: 29547635, <https://doi.org/10.1371/journal.pone.0194700>.

Celiac Disease Foundation. 2017. "Arsenic in the Gluten-Free Diet: Are You at Risk?" <https://celiac.org/blog/2017/05/arsenic-gluten-free-diet-risk/> [accessed 8 April 2018].

Children's Environmental Health and Disease Prevention Research Center at Dartmouth. 2016. "Arsenic." <http://www.dartmouth.edu/~childrenshealth/arsenic/> [accessed 23 March 2018].

Codex Alimentarius. 2017. "Code of Practice for the Prevention and Reduction of Arsenic Contamination in Rice." http://www.fao.org/fao-who-codexalimentarius/sh-proxy/en/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252Fstandards%252FCAC%252FBRC%252B77-2017%252FCXC_077e.pdf [accessed 18 April 2018].

Consumer Reports. 2015. "How Much Arsenic Is in Your Rice?" <https://www.consumerreports.org/cro/magazine/2015/01/how-much-arsenic-is-in-your-rice/index.htm> [accessed 23 March 2018].

Cubadda F, D'Amato M, Aureli F, Raggi A, Mantovani A. 2016. Dietary exposure of the Italian population to inorganic arsenic: The 2012–2014 Total Diet Study. *Food Chem Toxicol* 98(Pt B):148–158, PMID: 27756704, <https://doi.org/10.1016/j.fct.2016.10.015>.

Cubadda F, Jackson BP, Cottingham KL, Van Horne YO, Kurzius-Spencer M. 2017. Human exposure to dietary inorganic arsenic and other arsenic species: state of knowledge, gaps and uncertainties. *Sci Total Environ* 579:1228–1239, PMID: 27914647, <https://doi.org/10.1016/j.scitotenv.2016.11.108>.

Dartmouth College Superfund Research Program. 2017. "Arsenic and You." <http://www.arsenicandyou.org> [accessed 23 March 2018].

Davis MA, Li Z, Gilbert-Diamond D, Mackenzie TA, Cottingham KL, Jackson BP, et al. 2014. Infant toenails as a biomarker of in utero arsenic exposure. *J Expo Sci Environ Epidemiol* 24(5):467, PMID: 24896769, <https://doi.org/10.1038/jes.2014.38>.

Davis MA, Signes-Pastor AJ, Argos M, Slaughter F, Pendergrast C, Punshon T, et al. 2017. Assessment of human dietary exposure to arsenic through rice. *Sci Total Environ* 586:1237–1244, PMID: 28233618, <https://doi.org/10.1016/j.scitotenv.2017.02.119>.

Díaz O, Tapia Y, Muñoz O, Montoro R, Velez D, Almela C. 2012. Total and inorganic arsenic concentrations in different species of economically important algae harvested from coastal zones of Chile. *Food Chem Toxicol* 50(3–4):744–749, PMID: 22138359, <https://doi.org/10.1016/j.fct.2011.11.024>.

Food and Agriculture Organization. 2001. "Carrageenan." <http://www.fao.org/ag/agn/jecfa-additives/specs/Monograph1/Additive-117.pdf> [accessed 23 March 2018].

Gossai A, Zens MS, Punshon T, Jackson BP, Perry AE, Karagas MR. 2017. Rice consumption and squamous cell carcinoma of the skin in a United States population. *Environ Health Perspect* 125(9):097005, PMID: 28934722, <https://doi.org/10.1289/EHP1065>.

Hall M, Gamble M, Slavkovich V, Liu X, Levy D, Cheng Z, et al. 2007. Determinants of arsenic metabolism: blood arsenic metabolites, plasma folate, cobalamin, and homocysteine concentrations in maternal–newborn pairs. *Environ Health Perspect* 115(10):1503–1509, PMID: 17938743, <https://doi.org/10.1289/ehp.9906>.

Jackson BP, Taylor VF, Karagas MR, Punshon T, Cottingham KL. 2012. Arsenic, organic foods, and brown rice syrup. *Environ Health Perspect* 120(5):623–626, PMID: 227326149, <https://doi.org/10.1289/ehp.1104619>.

Jones MR, Tellez-Plaza M, Vaidya D, Grau M, Francesconi KA, Goessler W, et al. 2016. Estimation of inorganic arsenic exposure in populations with frequent seafood intake: evidence from MESA and NHANES. *Am J Epidemiol* 184(8):590–602, PMID: 27702745, <https://doi.org/10.1093/aje/kww097>.

Krisberg K. 2017. "Regulatory changes ignore science, threaten public health: new administration eroding progress." <http://thenationshealth.aphapublications.org/content/47/8/1.3> [accessed 16 March 2018].

Kurzius-Spencer M, Burgess JL, Harris RB, Hartz V, Roberge J, Huang S, et al. 2014. Contribution of diet to aggregate arsenic exposures—an analysis across populations. *J Expo Sci Environ Epidemiol* 24(2):156–162, PMID: 23860400, <https://doi.org/10.1038/jes.2013.37>.

Kurzius-Spencer M, da Silva V, Thomson CA, Hartz V, Hsu C-H, Burgess JL, et al. 2017. Nutrients in one-carbon metabolism and urinary arsenic methylation in the National Health and Nutrition Examination Survey (NHANES) 2003–2004. *Sci Total Environ* 607–608:381–390, PMID: 28697391, <https://doi.org/10.1016/j.scitotenv.2017.07.019>.

Lai PY, Cottingham KL, Steinmaus C, Karagas MR, Miller MD. 2015. Arsenic and rice: translating research to address health care providers' needs. *J Pediatr* 167(4):797–803, PMID: 26253210, <https://doi.org/10.1016/j.jpeds.2015.07.003>.

Li RY, Stroud JL, Ma JF, McGrath SP, Zhao FJ. 2009. Mitigation of arsenic accumulation in rice with water management and silicon fertilization. *Environ Sci Technol* 43(10):3778–3783, PMID: 19544887.

Limmer MA, Wise P, Dykes GE, Seyffert AL. 2018. Silicon decreases dimethylarsinic acid concentration in rice grain and mitigates straighthead disorder. *Environ Sci Technol* 52(8):4809–4816, PMID: 29608840, <https://doi.org/10.1021/acs.est.8b00300>.

Lundberg Family Farms. 2016. "A Letter from the CEO." <http://www.lundberg.com/info/arsenic-in-food/> [accessed 23 March 2018].

Meharg AA, Lombi E, Williams PN, Scheckel KG, Feldmann J, Raab A, et al. 2008. Speciation and localization of arsenic in white and brown rice grains. *Environ Sci Technol* 42(4):1051–1057, PMID: 18351071.

Melkonian S, Argos M, Hall MN, Chen Y, Parvez F, Pierce B, et al. 2013. Urinary and dietary analysis of 18,470 Bangladeshis reveal a correlation of rice consumption with arsenic exposure and toxicity. *PLoS One* 8(11):e80691, PMID: 24260455, <https://doi.org/10.1371/journal.pone.0080691>.

- Muraki I, Wu H, Imamura F, Laden F, Rimm EB, Hu FB, et al. 2015. Rice consumption and risk of cardiovascular disease: results from a pooled analysis of 3 U.S. cohorts. *Am J Clin Nutr* 101(1):164–172, PMID: 2527760, <https://doi.org/10.3945/ajcn.114.087551>.
- Mushak P, McKinzie M, Olson E, Metrick A. 2000. Arsenic and old laws: a scientific and public health analysis of arsenic occurrence in drinking water, its health effects, and EPA's outdated arsenic tap water standard. In: *Arsenic and Old Laws: A Scientific and Public Health*. New York, NY:NRDC Publications Department.
- Nachman KE, Ginsberg GL, Miller MD, Murray CJ, Nigra AE, Pendergrast CB. 2017. Mitigating dietary arsenic exposure: current status in the United States and recommendations for an improved path forward. *Sci Total Environ* 581:582:221–236, PMID: 28065543, <https://doi.org/10.1016/j.scitotenv.2016.12.112>.
- Nature's One. 2018. "Nature Knows What's Best for Your Baby." <https://www.naturesone.com/our-story/>, [accessed 23 March 2018].
- Naujokas MF, Anderson B, Ahsan H, Aposhian HV, Graziano JH, Thompson C, et al. 2013. The Broad scope of health effects from chronic arsenic exposure: update on a worldwide public health problem. *Environ Health Perspect* 121(3):295–302, PMID: 23458756, <https://doi.org/10.1289/ehp.1205875>.
- Nigra AE, Sanchez TR, Nachman KE, Harvey DE, Chillrud SN, Graziano JH, et al. 2017. The effect of the Environmental Protection Agency maximum contaminant level on arsenic exposure in the USA from 2003 to 2014: an analysis of the National Health and Nutrition Examination Survey (NHANES). *Lancet Public Health* 2(11):e513–e521, PMID: 29250608, [https://doi.org/10.1016/S2468-2667\(17\)30195-0](https://doi.org/10.1016/S2468-2667(17)30195-0).
- Norton GJ, Islam MR, Deacon CM, Zhao F-J, Stroud JL, McGrath SP, et al. 2009. Identification of low inorganic and total grain arsenic rice cultivars from Bangladesh. *Environ Sci Technol* 43(15):6070–6075, PMID: 19731720.
- Norton GJ, Pinson SR, Alexander J, Mckay S, Hansen H, Duan GL, et al. 2012. Variation in grain arsenic assessed in a diverse panel of rice (*Oryza sativa*) grown in multiple sites. *New Phytol* 193(3):650–664, PMID: 22142234, <https://doi.org/10.1111/j.1469-8137.2011.03983.x>.
- NRC (National Research Council). 2014. "Critical Aspects of EPA's IRIS Assessment of Inorganic Arsenic: Interim Report." Washington, DC: National Academies Press. <http://www.nap.edu/catalog/18594/critical-aspects-of-epas-iris-assessment-of-inorganic-arsenic-interim> [accessed 3 August 2018].
- NRC. 2018. "Progress toward Transforming the Integrated Risk Information System (IRIS) Program: A 2018 Evaluation (2018)." Washington, DC: National Academies Press. <https://www.nap.edu/download/25086> [accessed 18 April 2018].
- Penido ES, Bennett AJ, Hanson TE, Seyfferth AL. 2016. Biogeochemical impacts of silicon-rich rice residue incorporation into flooded soils: implications for rice nutrition and cycling of arsenic. *Plant Soil* 399(1–2):75–87, <https://doi.org/10.1007/s11104-015-2682-3>.
- Punshon T, Jackson BP, Meharg AA, Warczack T, Scheckel K, Guerinet ML. 2017. Understanding arsenic dynamics in agronomic systems to predict and prevent uptake by crop plants. *Sci Total Environ* 581–582:209–220, PMID: 28043702, <https://doi.org/10.1016/j.scitotenv.2016.12.111>.
- Punshon T, Jackson B. 2018. Essential micronutrient and toxic trace element concentrations in gluten containing and gluten-free foods. *Food Chem* 252:258–264, PMID: 29478539, <https://doi.org/10.1016/j.foodchem.2018.01.120>.
- Raab A, Baskaran C, Feldmann J, Meharg AA. 2009. Cooking rice in a high water to rice ratio reduces inorganic arsenic content. *J Environ Monit* 11(1):41–44, PMID: 19137137, <https://doi.org/10.1039/B816906C>.
- Samet JM, Burke TA, Goldstein BD. 2017. The Trump administration and the environment — heed the science. *N Engl J Med* 376(12):1182–1188, PMID: 28249122, <https://doi.org/10.1056/NEJMms1615242>.
- Schoof R, Yost L, Eickhoff J, Creelius E, Cragin D, Meacher D, et al. 1999. A market basket survey of inorganic arsenic in food. *Food Chem Toxicol* 37(8):839–846, PMID: 10506007.
- Sengupta M, Hossain M, Mukherjee A, Ahamed S, Das B, Nayak B, et al. 2006. Arsenic burden of cooked rice: traditional and modern methods. *Food Chem Toxicol* 44(11):1823–1829, PMID: 16876928, <https://doi.org/10.1016/j.fct.2006.06.003>.
- Seyfferth AL, Morris AH, Gill R, Kearns KA, Mann JN, Paukett M, et al. 2016. Soil incorporation of silica-rich rice husk decreases inorganic arsenic in rice grain. *J Agric Food Chem* 64(19):3760–3766, PMID: 27109244, <https://doi.org/10.1021/acs.jafc.6b01201>.
- Signes-Pastor AJ, Carey M, Meharg AA. 2017a. Inorganic arsenic removal in rice bran by percolating cooking water. *Food Chem* 234:76–80, PMID: 28551270, <https://doi.org/10.1016/j.foodchem.2017.04.140>.
- Signes-Pastor AJ, Woodside JV, McMullan P, Mullan K, Carey M, Karagas MR, et al. 2017b. Levels of infants' urinary arsenic metabolites related to formula feeding and weaning with rice products exceeding the EU inorganic arsenic standard. *PLoS One* 12(5):e0176923, PMID: 28472079, <https://doi.org/10.1371/journal.pone.0176923>.
- Signes-Pastor AJ, Cottingham KL, Carey M, Sayarath V, Palys T, Meharg AA, et al. 2018. Infants' dietary arsenic exposure during transition to solid food. *Sci Rep* 8(1):7114, PMID: 29739998, <https://doi.org/10.1038/s41598-018-25372-1>.
- Signes A, Mitra K, Burló F, Carbonell-Barrachina A. 2008. Effect of two different rice dehusking procedures on total arsenic concentration in rice. *Eur Food Res Technol* 226(3):561–567, <https://doi.org/10.1007/s00217-007-0571-6>.
- Spratlen MJ, Gamble MV, Grau-Perez M, Kuo C-C, Best LG, Yracheta J, et al. 2017. Arsenic metabolism and one-carbon metabolism at low-moderate arsenic exposure: evidence from the Strong Heart Study. *Food Chem Toxicol* 105:387–397, PMID: 28479390, <https://doi.org/10.1016/j.fct.2017.05.004>.
- Taylor V, Goodale B, Raab A, Schwerdtle T, Reimer K, Conklin S, et al. 2017. Human exposure to organic arsenic species from seafood. *Sci Total Environ* 580:266–282, PMID: 28024743, <https://doi.org/10.1016/j.scitotenv.2016.12.113>.
- Teasley WA, Limmer MA, Seyfferth AL. 2017. How rice (*Oryza sativa* L.) responds to elevated as under different Si-rich soil amendments. *Environ Sci Technol* 51(18):10335–10343, PMID: 28795805, <https://doi.org/10.1021/acs.est.7b01740>.
- The National Academies. 2018. "The Environmental Health Matters Initiative." <http://nas-sites.org/envirohealthmatters/background/> [accessed 30 April 2018].
- U.S. Congress. 2012. H.R.3984 – Apple Juice Act of 2012. <https://www.congress.gov/bills/112th-congress/house-bill/3984> [accessed 16 March 2018].
- U.S. Congress. 2015. H.R.2529 – Reducing Food-Based Inorganic Compounds Exposure Act of 2015. <https://www.congress.gov/bills/114th-congress/house-bill/2529/text> [accessed 16 March 2018].
- U.S. Congress. 2017. H.R.4535 – Rice Act. <https://www.congress.gov/bills/115th-congress/house-bill/4535> [accessed 18 April 2018].
- USDA. (U.S. Department of Agriculture). 2018. WVEIA/NHANES Overview. <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/wveianhanes-overview/> [accessed 4 May 2018].
- U.S. EPA (U.S. Environmental Protection Agency). 2000. "Arsenic in Drinking Water Rule Economic Analysis (EPA 815-R-00-026)." <https://nepis.epa.gov/Exe/ZyPdf.cgi?Dockey=20001YQT.txt> [accessed 9 July 2018].
- U.S. FDA (U.S. Food and Drug Administration). 2013. "Arsenic in Apple Juice." <https://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm280209.htm> [accessed 15 March 2018].
- U.S. FDA. 2016. "Arsenic in Rice and Rice Products." <https://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm319870.htm> [accessed 15 March 2018].
- U.S. FDA. 2017a. "Arsenic." <https://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm280202.htm> [accessed 23 March 2018].
- U.S. FDA. 2017b. "Combination Metals Testing." <https://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm521427.htm> [accessed 23 March 2018].
- U.S. GAO (U.S. Government Accountability Office). 2018. "Federal Efforts to Manage the Risk of Arsenic in Rice." <https://www.gao.gov/assets/700/690701.pdf> [accessed 18 April 2018].
- Welch B, Smit E, Cardenas A, Hystad P, Kile ML. 2018. Trends in urinary arsenic among the U.S. population by drinking water source: results from the National Health and Nutritional Examinations Survey 2003–2014. *Environ Res* 162:8–17, PMID: 29272814, <https://doi.org/10.1016/j.envres.2017.12.012>.
- Wright RO, Christiani D. 2010. Gene-environment interaction and children's health and development. *Curr Opin Pediatr* 22(2):197, PMID: 20090521, <https://doi.org/10.1097/MOP.0b013e328336ebf9>.
- Xue J, Zartarian V, Wang S-W, Liu SV, Georgopoulos P. 2010. Probabilistic modeling of dietary arsenic exposure and dose and evaluation with 2003–2004 NHANES data. *Environ Health Perspect* 118(3):345, PMID: 20194069, <https://doi.org/10.1289/ehp.0901205>.
- Yang J, Zhou Q, Zhang J. 2017. Moderate wetting and drying increases rice yield and reduces water use, grain arsenic level, and methane emission. *Crop J* 5(2):151–158, <https://doi.org/10.1016/j.cj.2016.06.002>.
- Zhang R, Zhang X, Wu K, Wu H, Sun Q, Hu FB, et al. 2016. Rice consumption and cancer incidence in US men and women. *Int J Cancer* 138(3):555–564, PMID: 26219234, <https://doi.org/10.1002/ijc.29704>.